

Class IX Session 2024-25

Subject - Mathematics

Sample Question Paper - 5

Time Allowed: 3 hours

Maximum Marks: 80

General Instructions:


1. This Question Paper has 5 Sections A-E.
2. Section A has 20 MCQs carrying 1 mark each.
3. Section B has 5 questions carrying 02 marks each.
4. Section C has 6 questions carrying 03 marks each.
5. Section D has 4 questions carrying 05 marks each.
6. Section E has 3 case based integrated units of assessment carrying 04 marks each.
7. All Questions are compulsory. However, an internal choice in 2 Qs of 5 marks, 2 Qs of 3 marks and 2 Questions of 2 marks has been provided. An internal choice has been provided in the 2 marks questions of Section E.
8. Draw neat figures wherever required. Take $\pi = 22/7$ wherever required if not stated.

Section A


1. The point which lies on y-axis at a distance of 6 units in the positive direction of y-axis is [1]

a) (-6, 0)	b) (0, -6)
c) (6, 0)	d) (0, 6)
2. The perimeter of an equilateral triangle is 60 m. The area is [1]

a) $10\sqrt{3} m^2$	b) $20\sqrt{3} m^2$
c) $15\sqrt{3} m^2$	d) $100\sqrt{3} m^2$
3. In a figure, O is the centre of the circle with AB as diameter. If $\angle AOC = 40^\circ$, the value of x is equal to [1]



- | | |
|---------------|---------------|
| a) 80° | b) 50° |
| c) 70° | d) 60° |
4. In the given figure, ABCD is a parallelogram in which $\angle BDC = 45^\circ$ and $\angle BAD = 75^\circ$. Then, $\angle CBD = ?$ [1]




- a) 60°
- b) 45°
- c) 75°
- d) 55°

5. The product $\sqrt[3]{2} \cdot \sqrt[4]{2} \cdot \sqrt[12]{32}$ is equal to [1]

- a) $\sqrt[12]{2}$
- b) 2
- c) $\sqrt{2}$
- d) $\sqrt[12]{32}$

6. In the given figure, the sides CB and BA of $\triangle ABC$ have been produced to D and E respectively such that $\angle ABD = 110^\circ$ and $\angle CAE = 135^\circ$. Then, $\angle ACB = ?$ [1]



- a) 35°
- b) 45°
- c) 65°
- d) 55°

7. If $x = 3$ and $y = -2$ satisfies $5x - y = k$, then the value of k is [1]

- a) 3
- b) 17
- c) 12
- d) -2

8. The degree of the zero polynomial is [1]

- a) 0
- b) any natural number
- c) 1
- d) not defined

9. The decimal form of $\frac{2}{11}$ is [1]

- a) 0.018
- b) 0.18
- c) $0.\overline{18}$
- d) $0.0\overline{18}$

10. If one angle of a parallelogram is 24° less than twice the smallest angle, then the measure of the largest angle of the parallelogram is [1]


- a) 112°
- b) 68°
- c) 176°
- d) 102°

11. $9^3 + (-3)^3 - 6^3 = ?$ [1]

- a) 540
c) 270
- b) 486
d) 432

12. The equation $x - 2 = 0$ on number line is represented by [1]
 a) infinitely many lines
b) two lines
c) a point
d) a line

13. In the given figure, $\angle OAB = 110^\circ$ and $\angle BCD = 130^\circ$ then $\angle ABC$ is equal to [1]




- a) 50°
c) 40°
- b) 60°
d) 70°

14. If $\frac{5-\sqrt{3}}{2+\sqrt{3}} = x + y\sqrt{3}$, then [1]

- a) $x = -13, y = -7$
c) $x = -13, y = 7$
- b) $x = 13, y = -7$
d) $x = 13, y = 7$

15. In Fig. ABCD is a cyclic quadrilateral. If $\angle BAC = 50^\circ$ and $\angle DBC = 60^\circ$ then find $\angle BCD$. [1]



- a) 50°
c) 70°
- b) 60°
d) 55°

16. Which of the following points lies on the line $y = 2x + 3$? [1]

- a) (2,8)
c) (3,9)
- b) (5,15)
d) (4,12)

17. How many lines pass through two points? [1]

- a) many
c) two
- b) three
d) only one

18. Which one of the following is a polynomial? [1]

- a) $\frac{x-1}{x+1}$
c) $x^2 + \frac{3x^2}{\sqrt{x}}$
- b) $\sqrt{2x} - 1$
d) $\frac{x^2}{2} - \frac{2}{x^2}$

19. **Assertion (A):** In $\triangle ABC$, median AD is produced to X such that $AD = DX$. Then ABXC is a parallelogram. [1]

Reason (R): Diagonals AX and BC bisect each other at right angles.

- a) Both A and R are true and R is the correct explanation of A. b) Both A and R are true but R is not the correct explanation of A.
- c) A is true but R is false. d) A is false but R is true.

20. **Assertion (A):** Three rational numbers between $\frac{2}{5}$ and $\frac{3}{5}$ are $\frac{9}{20}$, $\frac{10}{20}$ and $\frac{11}{20}$ [1]

Reason (B): A rational number between two rational numbers p and q is $\frac{1}{2}(p+q)$


- a) Both A and R are true and R is the correct explanation of A. b) Both A and R are true but R is not the correct explanation of A.
- c) A is true but R is false. d) A is false but R is true.

Section B

21. If a point O lies between two points P and R such that $PO = OR$ then prove that $PO = \frac{1}{2}PR$. [2]

22. Why is Axiom 5, in the list of Euclid's axioms, considered a **universal truth**? [2]

23. In Fig., if ABC and ABD are equilateral triangles then find the coordinates of C and D. [2]



24. Prove that: $\frac{a+b+c}{a^{-1}b^{-1}+b^{-1}c^{-1}+c^{-1}a^{-1}} = abc$ [2]

OR

Express $0.35\bar{7}$ in the form $\frac{p}{q}$, where p and q are integers and $q \neq 0$.

25. If the volume of a right circular cone of height 9 cm is $48\pi \text{ cm}^3$, find the diameter of its base. [2]

OR

A team of 10 interns and 1 professor from zoological department visited a forest, where they set up a conical tent for their accommodation. There they perform activities like planting saplings, yoga, cleaning lakes, testing the water for contaminants and pollutant levels and desilt the lake bed and also using the silt to strengthen bunds.

Find the radius and height of the tent if the base area of tent is 154 cm^2 and curved surface area of the tent is 396 cm^2 .

Section C

26. Represent $\sqrt{4.5}$ on the number line. [3]

27. Draw a histogram for the daily earnings of 30 drug stores in the following table: [3]

Daily earnings (in ₹):	450 - 500	500 - 550	550 - 600	600 - 650	650 - 700
Number of Stores:	16	10	7	3	1

28. ABC is a triangle right angled at C. A line through the mid-point M of hypotenuse AB and parallel to BC [3]

intersects AC at D. Show that

- i. D is the mid-point of AC
- ii. $MD \perp AC$
- iii. $CM = MA = \frac{1}{2}AB$

29. Write linear equation $3x + 2y = 18$ in the form of $ax + by + c = 0$. Also write the values of a, b and c. Are (4, 3) and (1, 2) solution of this equation? [3]

30. Following are the marks of a group of 92 students in a test of reading ability : [3]

Marks	50-52	47-49	44-46	41-43	38-40	35-37	32-34	Total
Number of students	4	10	15	18	20	12	13	92

Construct a frequency polygon for the above data.

OR


Draw a frequency polygon for the following distribution:

Marks obtained	0-10	10-20	20-30	30-40	40-50	50-60	60-70	70-80
No. of students	7	10	6	8	12	3	2	2

31. The polynomials $ax^3 + 3x^2 - 3$ and $2x^3 - 5x + a$ when divided by $(x - 4)$ leave the remainders R_1 and R_2 respectively. Find the values of a if $R_1 + R_2 = 0$ [3]


Section D

32. In each of the figures given below, $AB \parallel CD$. Find the value of x° in each case. [5]



OR

In the given figure, $AB \parallel CD$ and $\angle AOC = x^\circ$. If $\angle OAB = 104^\circ$ and $\angle OCD = 116^\circ$, find the value of x.



33. An iron pillar consists of a cylindrical portion 2.8 m high and 20 cm in diameter and a cone 42 cm high is surmounting it. Find the weight of the pillar, given that 1 cm^3 of iron weighs 7.5 g. [5]

34. Find the percentage increase in the area of a triangle if its each side is doubled. [5]

OR

The sides of a triangle are in the ratio 5 : 12 : 13 and its perimeter is 150 m. Find the area of the triangle.

35. Find the integral roots of the polynomial $f(x) = x^3 + 6x^2 + 11x + 6$. [5]

Section E

36. Read the following text carefully and answer the questions that follow: [4]


Ajay lives in Delhi, The city of Ajay's father in laws residence is at Jaipur is 600 km from Delhi. Ajay used to travel this 600 km partly by train and partly by car.

He used to buy cheap items from Delhi and sale at Jaipur and also buying cheap items from Jaipur and sale at

Delhi.

Once From **Delhi to Jaipur** in forward journey he covered $2x$ km by train and the rest y km by taxi.

But, while returning he did not get a reservation from Jaipur in the train. So first $2y$ km he had to travel by taxi and the rest x km by Train. From Delhi to Jaipur he took 8 hrs but in returning it took 10 hrs.



- Write the above information in terms of equation. (1)
- Find the value of x and y ? (1)
- Find the speed of Taxi? (2)

OR


Find the speed of Train? (2)

37. Read the following text carefully and answer the questions that follow:

[4]

In the middle of the city, there was a park ABCD in the form of a parallelogram form so that $AB = CD$, $AB \parallel CD$ and $AD = BC$, $AD \parallel BC$.

Municipality converted this park into a rectangular form by adding land in the form of $\triangle APD$ and $\triangle BCQ$. Both the triangular shape of land were covered by planting flower plants.




- Show that $\triangle APD$ and $\triangle BQC$ are congruent. (1)
- PD is equal to which side? (1)
- Show that $\triangle ABC$ and $\triangle CDA$ are congruent. (2)

OR

What is the value of $\angle m$? (2)

38. Two new roads, Road E and Road F were constructed between society 4 and 1 and society 1 and 2.

[4]



- What would be the measure of the sum of angles formed by the straight roads at Society 1 and society 3?

- a. 60°
 - b. 90°
 - c. 180°
 - d. 360°
- ii. Krish says, The distance to go from society 4 to society 2 using Road D will be longer than the distance using Road E. Is Krish correct? Justify your answer with examples.
- iii. Road G, perpendicular to Road F was constructed to connect the park and Road F. Which of the following is true for Road G and Road F?
- a. Road G and road F are of same length.
 - b. Road F divides Road G into two equal parts.
 - c. Road G divides Road F into two equal parts.
 - d. The length of road G is one-fourth of the length of Road F.
- iv. Priya said, Minor arc corresponding to Road B is congruent to minor arc corresponding to Road D. Do you agree with Priya? Give reason to support your answer.





Today is your
OPPORTUNITY to build the
TOMORROW you want.

ADMISSIONS OPEN

Session 2024-25

for

AMU XI ENTRANCE

Science / Diploma / Commerce / Humanities

Batches Starting Soon

Vineet Coaching & Guidance Centre

VCGC Tower, Phase I, ADA Colony, Ramghat Road, Aligarh-202001 (UP)
website : www.vcgc.in | E-Mail : vcgc.official@gmail.com

 **8923803150, 9997447700**

XI AMU ENTRANCE RESULTS 2023-24



Aakash Gaur



Kanika Garg



Riddhima Tomar



Tanmay Mudgal



Aastha Sharma



Ayush Senger



Prabhat Singh



Anubhav Gupta



Akash Basu



Shaurya Gangal



Abhishek Gaur



Rudraksh Chaudhary



Aryan Tiwari



Pakhi Garg



Rashi Singh



Ragini Singh



Pratishtha Sharma



Aaliya Singh



Kanika Singh



Yashika



Manav Kushwaha



Saloni Chaudhary



Ayushi Dhanger



Ayushi Gautam



Mugdha Singh



Afifa Malik



Khushi Varshney



Bhoomi Agarwal



Sanchi Popli



Shubhi Awasthi



Prisha Tanwar



Khanak



Sanyogita



Aakashi Sharma



Ipshita Upadhyay



Yashi Vashishtha



Divya Singh




Karnika Singh



Harshit Chaudhary



Vivek Gautam



Lalit Dhargar



Prachi Singh



Harshit Raghav



Srishty



Vanshika Garg



Aman Varshney



Pranjal Tiwari



Priyanshi Dhanger



Purnank Nandan

and many more...

 **8923803150, 9997447700**



@vcgc.aligarh



@vcgc_aligarh



@VCGC Aligarh



VCGC Online



VCGC Online App

Solution

Section A

1.

(d) (0, 6)

Explanation: Since it lies on the y-axis so its abscissa x will be zero.
Thus, the point will be (0, 6).

2.

(d) $100\sqrt{3} m^2$

Explanation: Perimeter of equilateral triangle = 60 m

$$\Rightarrow 3 \times \text{side} = 60 \text{ m}$$

$$\Rightarrow \text{side} = 20 \text{ m}$$

$$\text{Area of equilateral triangle} = \frac{\sqrt{3}}{4}(\text{Side})^2$$


$$= \frac{\sqrt{3}}{4} 20 \times 20$$

$$= 100\sqrt{3} \text{ sq.m}$$

3.

(c) 70°

Explanation:



OA = OC (radii)

So, $\angle OAC = \angle OCA = x$

Again, In $\triangle OAC$

$$\angle OAC + \angle OCA + \angle AOC = 180^\circ$$

$$x + x + \angle AOC = 180^\circ$$

$$x + x + 40^\circ = 180^\circ$$

$$2x = 140^\circ$$

$$x = 70^\circ$$

4. (a) 60°

Explanation: As per the question

$$\angle BAD = \angle BCD = 75^\circ \text{ (opposite angles of parallelogram)}$$

Now, in $\triangle BCD$,

$$\angle BCD + \angle CBD + \angle BDC = 180^\circ$$

$$45^\circ + \angle CBD + 75^\circ = 180^\circ$$

$$\angle CBD = 60^\circ$$

5.

(b) 2

Explanation: $\sqrt[3]{2} \cdot \sqrt[4]{2} \cdot \sqrt[12]{32}$

$$= \sqrt[3]{2} \cdot \sqrt[4]{2} \cdot \sqrt[12]{(2)^5}$$

$$= (2)^{\frac{1}{3}} \cdot (2)^{\frac{1}{4}} \cdot (2)^{\frac{5}{12}}$$



$$= (2)^{\frac{1}{3} + \frac{1}{4} + \frac{5}{12}}$$

$$= (2)^{\frac{4+3+5}{12}}$$

$$= (2)^{\frac{12}{12}}$$

$$= 2$$

6.

(c) 65°

Explanation: We can find $\angle CBA$ as follows:

Given that $\angle EBA = 110^\circ$

$\angle CBA = 180 - 110 \dots$ (linear pair)

$= 70^\circ$

Given $\angle CAD = 135^\circ$

So, $\angle CAB = 180 - 135 \dots$ (linear pair)

$= 45^\circ$

So, $\angle ACB = 180 - (70 + 45) \dots$ (angle sum property of triangle)

$= 65^\circ$

7.

(b) 17

Explanation: If $x = 3$ and $y = -2$ satisfies $5x - y = k$

Then

$5x - y = k$

$5 \times 3 - (-2) = k$

$15 + 2 = k$

$k = 17$

8.

(d) not defined

Explanation: The general form of a polynomial is $a_n x^n$, where n is a natural number.

For zero polynomial $a_n = 0$.

Since the largest value of n for which a_n is non-zero is negative infinity (all the integers are bigger than negative infinity).

Therefore, the degree of zero polynomials is not defined.

9.

(c) $0.\overline{18}$

Explanation: When we divide 2 by 11

We have value $= 0.181818\dots$


Which is $0.\overline{18}$

10.

(a) 112°

Explanation:

Let angles of parallelogram are $\angle A, \angle B, \angle C, \angle D$



Let smallest angle $= \angle A$

Let largest angle $= \angle B$

$= \angle B = 2\angle A - 24^\circ \dots$ (i)

$\angle A + \angle B = 180^\circ$ [adjacent angle of parallelogram]

So, $\angle A + 2\angle A - 24^\circ = 180^\circ$

$= 3\angle A = 180^\circ + 24^\circ = 204^\circ$

$$= \angle A = \frac{204^\circ}{3} = 68^\circ$$

$$= \angle B = 2 \times 68^\circ - 24^\circ = 112^\circ$$

11.

(b) 486

Explanation: $9^3 + (-3)^3 - 6^3$

$$= 729 - 27 - 216$$

$$= 729 - 243$$

$$= 486$$

12.

(c) a point

Explanation: $x - 2 = 0$

$x = 2$ is a point on the number line


13.

(b) 60°

Explanation:

In the given figure, $OA \parallel CD$.

Construction: Extend OA such that it intersects BC at E.



Now, $OE \parallel CD$ and BC is a transversal.

$\therefore \angle AEC = \angle BCD = 130^\circ$ (Pair of corresponding angles)

Also, $\angle OAB + \angle BAE = 180^\circ$ (Linear pair)

$$\therefore 110^\circ + \angle BAE = 180^\circ$$

$$\Rightarrow \angle BAE = 180^\circ - 110^\circ = 70^\circ$$

In $\triangle ABE$

$\angle AEC = \angle BAE + \angle ABE$... (In a triangle, exterior angle is equal to the sum of two opposite interior angles)

$$\therefore 130^\circ = 70^\circ + x^\circ$$

$$\Rightarrow x^\circ = 130^\circ - 70^\circ = 60^\circ$$

Thus, the measure of angle $\angle ABC$ is 60°

14.

(b) $x = 13, y = -7$

Explanation: $x + y\sqrt{3} = \frac{5-\sqrt{3}}{2+\sqrt{3}}$

$$= \frac{5-\sqrt{3}}{2+\sqrt{3}} \times \frac{2-\sqrt{3}}{2-\sqrt{3}}$$

$$= \frac{(5-\sqrt{3})(2-\sqrt{3})}{(2)^2 - (\sqrt{3})^2}$$

$$= \frac{5(2-\sqrt{3}) - \sqrt{3}(2-\sqrt{3})}{4-3}$$

$$= \frac{10-5\sqrt{3}-2\sqrt{3}+3}{1}$$

$$= 13 - 7\sqrt{3}$$


$$\text{Hence, } x + y\sqrt{3} = 13 - 7\sqrt{3}$$

$$\Rightarrow x = 13, y = -7$$

15.

(c) 70°

Explanation: Here $\angle BDC = \angle BAC = 50^\circ$ (angles in same segment are equal)



In $\triangle BCD$, we have
 $\angle BCD = 180^\circ - (\angle BDC + \angle DBC)$
 $= 180^\circ - (50^\circ + 60^\circ)$
 $= 70^\circ$

16.

(c) (3,9)

Explanation: Here, $y = 2x + 3$

So, for $x = 3$, we have

$$y = 2 \times 3 + 3$$

$$= 6 + 3$$

$$= 9$$

So, (3, 9) lies on the given line

17.

(d) only one

Explanation: only one because if a line is passing through two points then that two points are solution of a single linear equation

so only one line passes over two given points.

18.

(c) $x^2 + \frac{3x^{\frac{3}{2}}}{\sqrt{x}}$

Explanation: Since the power of the variable of all terms of a polynomial should be a whole number. Then

$$x^2 + \frac{3x^{\frac{3}{2}}}{\sqrt{x}}$$

$$= x^2 + 3x^{\frac{3}{2} - \frac{1}{2}}$$

$$= x^2 + 3x^{\frac{2}{2}}$$

$$= x^2 + 3x$$

Here the powers of variable are whole numbers. Therefore the given expression is a polynomial.

19.


(c) A is true but R is false.

Explanation:

In quadrilateral ABXC, we have

$$AD = DX \text{ [Given]}$$

$$BD = DC \text{ [Given]}$$



So, diagonals AX and BC bisect each other but not at right angles.

Therefore, ABXC is a parallelogram.

20. (a) Both A and R are true and R is the correct explanation of A.

Explanation: Both A and R are true and R is the correct explanation of A.

Section B



Given : In the fig. PR is a line segment such that, PO = OR,

Proof: From Fig. we have

$$PO + OR = PR \dots(i)$$

$$PO = OR \text{ (Given) } \dots(ii)$$

$$PO + PO = PR \text{ [Using (ii) in (i)]}$$

$$2PO = PR$$

$$\text{Therefore } PO = \frac{1}{2}PR$$

22. Euclid's Axiom 5 states that "The whole is greater than the part. Since this is true for anything in any part of the world. So, this is a universal truth.

23. Here, AC = 2a and AO = a

By Pythagoras theorem

$$OC^2 = AC^2 - AO^2 = 4a^2 - a^2 = 3a^2$$

$$OC = a\sqrt{3}$$

Therefore, coordinates of C are (0, $a\sqrt{3}$)

And the coordinates of D are (0, $-a\sqrt{3}$).

24. LHS

$$= \frac{a+b+c}{a^{-1}b^{-1}+b^{-1}c^{-1}+c^{-1}a^{-1}}$$

$$= \frac{a+b+c}{\frac{1}{a} \times \frac{1}{b} + \frac{1}{b} \times \frac{1}{c} + \frac{1}{c} \times \frac{1}{a}}$$

$$= \frac{a+b+c}{\frac{1}{ab} + \frac{1}{bc} + \frac{1}{ca}}$$

$$= \frac{a+b+c}{\frac{abc}{abc} + \frac{abc}{abc} + \frac{abc}{abc}}$$

$$= \frac{abc(a+b+c)}{abc}$$

$$= abc$$

$$= \text{RHS}$$

$$\text{LHS} = \text{RHS}$$

$$\text{Hence Proved}$$

Let $x = 0.35\overline{7}$.

Then, $x = 0.35777\dots$

$$\text{So, } 100x = 35.777\dots \dots(i)$$

$$1000x = 357.777\dots \dots(ii)$$

Subtracting (i) from (ii), we get

$$1000x - 100x = 357.777\dots - 35.777\dots$$

$$900x = 322$$

$$\Rightarrow x = \frac{322}{900}$$

$$\Rightarrow x = \frac{322}{900} = \frac{161}{450}$$

25. Let the radius of the base of the right circular cone be r cm.

$$h = 9 \text{ cm, volume} = 48\pi \text{ cm}^3$$

$$\Rightarrow \frac{1}{3}\pi r^2 h = 48\pi$$

$$\Rightarrow \frac{1}{3}r^2 h = 48$$

$$\Rightarrow \frac{1}{3} \times r^2 \times 9 = 48$$

$$\Rightarrow r^2 = \frac{48 \times 3}{9}$$

$$\Rightarrow r^2 = 16 \Rightarrow r = \sqrt{16} = 4 \text{ cm}$$

$$\Rightarrow 2r = 2(4) = 8 \text{ cm.}$$

\therefore the diameter of the base of the right circular cone is 8 cm.

OR

A tent is of conical shape. Thus,

$$\text{Base area} = \pi r^2 = 154 \text{ cm}^2$$



So, radius $r = 7$ cm

$$\text{Curved surface area} = \pi r l = 396 \text{ cm}^2$$

$$396 = 3.14 \times 7 \times l$$

$$\Rightarrow l = 18 \text{ cm}$$

$$\text{Now, height } h = \sqrt{l^2 - r^2} = \sqrt{18^2 - 7^2} = 16.5 \text{ cm}$$

Section C

26. Consider, $AB = 4.5$ units.


Extend AB upto point C such that $BC = 1$ unit.

$$\therefore AC = 4.5 + 1 = 5.5 \text{ units.}$$

Now mark O as the midpoint of AC .

With O as centre and radius OC draw a semicircle.

Draw perpendicular BD on AC which intersect the semicircle at D .



This length $BD = \sqrt{4.5}$ units.

To show BD on the number line, consider line ABC as number line with point B as zero.

Therefore, $BC = 1$ unit.

With B as centre and radius BD draw an arc which intersects number line ABC at E .


So, point E represents $\sqrt{4.5}$

$AB = 4.5$ units

$BC = 1$ unit


$$BD = BE = \sqrt{4.5} \text{ units}$$

27. A histogram for the daily earnings of 30 drug stores



28. Given: ABC is a triangle right angled at C . A line through the mid-point M of hypotenuse AB and parallels to BC intersects AC at

D .



To Prove:

i. D is the mid-point of AC (ii) $MD \perp AC$

$$\text{ii. } CM = MA = \frac{1}{2}AB$$

Proof :

i. In ACB ,

As M is the mid-point of AB and $MD \parallel BC$

$\therefore D$ is the mid-point of AC . . . [By converse of mid-point theorem]

ii. As $MD \parallel BC$ and AC intersects them

$$\angle ADM = \angle ACB \dots [\text{Corresponding angles}]$$

$$\text{But } \angle ACB = 90^\circ \dots [\text{Given}]$$

$$\therefore \angle ADM = 90^\circ \Rightarrow MD \perp AC$$

iii. Now $\angle ADM + \angle CDM = 180^\circ \dots [\text{Linear pair axiom}]$

$$\angle ADM = \angle CDM = 90^\circ$$

In $\triangle ADM$ and $\triangle CDM$

$$AD = CD \dots [\text{As } D \text{ is the mid-point of } AC]$$

$$\angle ADM = \angle CDM \dots [\text{Each } 90^\circ]$$

$$DM = DM \dots [\text{Common}]$$

$$\therefore \triangle ADM \cong \triangle CDM \dots [\text{By SAS rule}]$$

$$\therefore MA = MC \dots [\text{c.p.c.t.}]$$

But M is the mid-point of AB

$$\therefore MA = MB = \frac{1}{2} AB$$

$$\therefore MA = MC = \frac{1}{2} AB$$

$$\therefore CM = MA = \frac{1}{2} AB$$

29. We have the equation as $3x + 2y = 18$

In standard form

$$3x + 2y - 18 = 0$$

$$\text{Or } 3x + 2y + (-18) = 0$$

But standard linear equation is

$$ax + by + c = 0$$

On comparison we get, $a = 3$, $b = 2$, $c = -18$

If $(4, 3)$ lie on the line, i.e., solution of the equation $LHS = RHS$

$$\therefore 3(4) + 2(3) = 18$$

$$12 + 6 = 18$$

$$18 = 18$$

As $LHS = RHS$, Hence $(4, 3)$ is the solution of given equation.

Again for $(1, 2)$

$$3x + 2y = 18$$

$$\therefore 3(1) + 2(2) = 18$$

$$3 + 4 = 18$$

$$7 = 18$$


$$LHS \neq RHS$$

Hence $(1, 2)$ is not the solution of given equation.


Therefore $(4, 3)$ is the point where the equation of the line $3x + 2y = 18$ passes through where as the line for the equation $3x + 2y = 18$ does not pass through the point $(1, 2)$.

30. First, we shall make the distribution continuous. Then we have,

Marks	Number of students
31.5-34.5	13
34.5-37.5	12
37.5-40.5	20
40.5-43.5	18
43.5-46.5	15
46.5-49.5	10
49.5-52.5	4



OR



x_i	f_i	(x_i, f_i)
5	7	(5, 7)
15	10	(15, 10)
25	6	(25, 6)
35	8	(35, 8)
45	12	(45, 12)
55	3	(55, 3)
65	2	(65, 2)
75	2	(75, 2)

31. The given polynomials are,

$$f(x) = ax^3 + 3x^2 - 3$$

$$p(x) = 2x^3 - 5x + a$$

Let,

R_1 is the remainder when $f(x)$ is divided by $x - 4$

$$\Rightarrow R_1 = f(4)$$

$$\Rightarrow R_1 = a(4)^3 + 3(4)^2 - 3$$

$$= 64a + 48 - 3$$

$$= 64a + 45 \dots(1)$$

Now, let

R_2 is the remainder when $p(x)$ is divided by $x - 4$

$$\Rightarrow R_2 = p(4)$$

$$\Rightarrow R_2 = 2(4)^3 - 5(4) + a$$

$$= 128 - 20 + a$$

$$= 108 + a \dots (2)$$

Given, $R_1 + R_2 = 0$


$$\Rightarrow 64a + 45 + 108 + a = 0$$

$$\Rightarrow 65a + 153 = 0$$

$$\Rightarrow a = -\frac{153}{65}$$

This is the required value of a .

Section D



Draw $EF \parallel AB \parallel CD$

Now, $AB \parallel EF$ and BE is the transversal.

Then,

$\angle ABE = \angle BEF$ [Alternate Interior Angles]

$$\Rightarrow \angle BEF = 35^\circ$$

Again, $EF \parallel CD$ and DE is the transversal

Then,

$\angle DEF = \angle FED$

$$\Rightarrow \angle FED = 65^\circ$$

$$\therefore x^\circ = \angle BEF + \angle FED$$

$$x^\circ = 35^\circ + 65^\circ$$

$$x^\circ = 100^\circ$$

Through O draw $OE \parallel AB \parallel CD$

Then, $\angle AOE + \angle COE = x^\circ$

Now, $AB \parallel OE$ and AO is the transversal

$$\therefore \angle OAB + \angle AOE = 180^\circ$$

$$\Rightarrow 104^\circ + \angle AOE = 180^\circ$$

$$\Rightarrow \angle AOE = (180 - 104)^\circ = 76^\circ \dots(1)$$

Again, $CD \parallel OE$ and OC is the transversal


$$\therefore \angle COE + \angle OCD = 180^\circ$$

$$\Rightarrow \angle COE + 116^\circ = 180^\circ$$

$$\Rightarrow \angle COE = (180^\circ - 116^\circ) = 64^\circ \dots(2)$$

$$\therefore \angle AOC = \angle AOE + \angle COE = (76^\circ + 64^\circ) = 140^\circ \text{ [from (1) and (2)]}$$

Hence, $x^\circ = 140^\circ$



33. We are Given that,

An iron pillar consists of a cylindrical portion and a cone mounted on it.

The height of the cylindrical portion of the pillar, $H = 2.8 \text{ m} = 280 \text{ cm}$.

The height of the conical portion of the pillar, $h = 42$ cm..

The diameter of the cylindrical portion of the pillar = diameter of the circular base of cone = $D = 20$ cm.

The radius of the circular base of cylinder/ cone $r = \frac{D}{2} = 10$ cm.

Now, we have,

Volume of the pillar, $(V) =$ Volume of the cylindrical portion of pillar + volume of the conical portion of the pillar.

$$\Rightarrow V = \pi r^2 H + \frac{1}{3} \pi r^2 h$$

$$\Rightarrow V = \left(\frac{22}{7} \times 10^2 \times 280 + \frac{1}{3} \times \frac{22}{7} \times 10^2 \times 42 \right) \text{ cm}^3$$

$$\Rightarrow V = (22 \times 100 \times 40 + 22 \times 100 \times 2) \text{ cm}^3$$

$$\Rightarrow V = (88000 + 4400) \text{ cm}^3$$

$$\Rightarrow V = 92400 \text{ cm}^3$$

Hence, volume of iron pillar is 92400 cm^3

Given,

Weight of 1 cm^3 iron = 7.5 gm.

Hence, weight of 92400 cm^3 iron = 7.5×92400 gm.

$$= 693000 \text{ gm.}$$

$$= 693 \text{ Kg.}$$

Since, $1\text{Kg} = 1000$ gm.

Hence, the weight of iron pillar is 693 Kg.

34. Let a, b, c be the sides of the old triangle and s be its semi-perimeter. Then,

$$s = \frac{1}{2}(a + b + c)$$

The sides of the new triangle are $2a, 2b$ and $2c$.

Let s' be its semi-perimeter. Then,

$$s' = \frac{1}{2}(2a + 2b + 2c) = a + b + c = 2s$$

$$\Rightarrow s' = 2s$$

Let Δ and Δ' be the areas of the old and new triangles respectively. Then

$$\Delta = \sqrt{s(s-a)(s-b)(s-c)} \dots\dots(1)$$

and

$$\Delta' = \sqrt{s'(s'-2a)(s'-2b)(s'-2c)}$$

$$\Rightarrow \Delta' = \sqrt{2s(2s-2a)(2s-2b)(2s-2c)} \quad [\because s' = 2s]$$

$$\Rightarrow \Delta' = \sqrt{16s(s-a)(s-b)(s-c)}$$

$$\Rightarrow \Delta' = 4\sqrt{s(s-a)(s-b)(s-c)} = 4\Delta \quad [\text{from (1)}]$$

$$\therefore \text{Increase in the area of the triangle} = \Delta' - \Delta = 4\Delta - \Delta = 3\Delta$$

$$\text{Hence, percentage increase in area} = \left(\frac{3\Delta}{\Delta} \times 100 \right) = 300\%$$

OR

Given that the sides of a triangle are in the ratio $5: 12: 13$ and its perimeter is 150 m

Let the sides of the triangle be $5x$ m, $12x$ m and $13x$ m.

We know:

Perimeter = Sum of all sides

$$\text{or, } 150 = 5x + 12x + 13x$$

$$\text{or, } 30x = 150$$

$$\text{or, } x = 5$$

Thus, we obtain the sides of the triangle.

$$5 \times 5 = 25 \text{ m}$$

$$12 \times 5 = 60 \text{ m}$$

$$13 \times 5 = 65 \text{ m}$$

Now,

Let:

$$a = 25 \text{ m, } b = 60 \text{ m and } c = 65 \text{ m}$$

$$\therefore s = \frac{150}{2} = 75 \text{ m}$$

$$\Rightarrow s = 75 \text{ m}$$

By Heron's formula, we have

$$\begin{aligned}
 \text{Area of triangle} &= \sqrt{s(s-a)(s-b)(s-c)} \\
 &= \sqrt{75(75-25)(75-60)(75-65)} \\
 &= \sqrt{75 \times 50 \times 15 \times 10} \\
 &= \sqrt{15 \times 5 \times 5 \times 10 \times 15 \times 10} \\
 &= 15 \times 5 \times 10 \\
 &= 750 \text{ m}^2
 \end{aligned}$$

35. Given, that $f(x) = x^3 + 6x^2 + 11x + 6$

Clearly we can say that, the polynomial $f(x)$ with an integer coefficient and the highest degree term coefficient which is known as leading factor is 1.

So, the roots of $f(x)$ are limited to integer factor of 6, they are $\pm 1, \pm 2, \pm 3, \pm 6$

Let $x = -1$

$$\begin{aligned}
 f(-1) &= (-1)^3 + 6(-1)^2 + 11(-1) + 6 \\
 &= -1 + 6 - 11 + 6 \\
 &= 0
 \end{aligned}$$

Let $x = -2$

$$\begin{aligned}
 f(-2) &= (-2)^3 + 6(-2)^2 + 11(-2) + 6 \\
 &= -8 - (6 \times 4) - 22 + 6 \\
 &= -8 + 24 - 22 + 6 \\
 &= 0
 \end{aligned}$$

Let $x = -3$

$$\begin{aligned}
 f(-3) &= (-3)^3 + 6(-3)^2 + 11(-3) + 6 \\
 &= -27 - (6 \times 9) - 33 + 6 \\
 &= -27 + 54 - 33 + 6 \\
 &= 0
 \end{aligned}$$

But from all the given factors only $-1, -2, -3$ gives the result as zero. Further, since $f(x)$ is a polynomial of degree 3, therefore, it has almost 3 roots.

Therefore, the integral roots of $x^3 + 6x^2 + 11x + 6$ are $-1, -2, -3$.

Section E

36. i. Delhi to Jaipur: $2x + y = 600$

Jaipur to Delhi: $2y + x = 600$

Let S_1 and S_2 be the speeds of Train and Taxi respectively, then

Delhi to Jaipur: $\frac{2x}{S_1} + \frac{y}{S_2} = 8 \dots(i)$

Jaipur to Delhi: $\frac{x}{S_1} + \frac{2y}{S_2} = 10 \dots(ii)$

ii. $2x + y = 600 \dots(1)$

$x + 2y = 600 \dots(2)$

Solving (1) and (2) $\times 2$

$$2x + y - 2x - 4y = 600 - 1200$$

$$\Rightarrow -3y = -600$$

$$\Rightarrow y = 200$$

Put $y = 200$ in (1)

$$2x + 200 = 600$$

$$\Rightarrow x = \frac{400}{2} = 200$$

iii. We know that speed = $\frac{\text{Distance}}{\text{Time}} \Rightarrow \text{Time} = \frac{\text{Distance}}{\text{Speed}}$

Let S_1 and S_2 are speeds of train and taxi respectively.

Delhi to Jaipur: $\frac{2x}{S_1} + \frac{y}{S_2} = 8 \dots(i)$

Jaipur to Delhi: $\frac{x}{S_1} + \frac{2y}{S_2} = 10 \dots(ii)$

Solving (i) and (ii) $\times 2$

$$\Rightarrow \frac{2x}{S_1} + \frac{y}{S_2} - \frac{2x}{S_1} - \frac{4y}{S_2} = 8 - 20 = -12$$

$$\Rightarrow \frac{-3y}{S_2} = -12$$

We know that $y = 200$ km

$$\Rightarrow S_2 = \frac{3 \times 200}{12} = 50 \text{ km/hr}$$

Hence speed of Taxi = 50 km/hr

OR

We know that $x = 200$ km

Put $S_2 = 50$ km/hr ... (i)

$$\frac{400}{S_1} + \frac{200}{50} = 8$$

$$\Rightarrow \frac{400}{S_1} = 8 - 4 = 4$$

$$\Rightarrow S_1 = \frac{400}{4} = 100 \text{ km/hr}$$

Hence speed of Train = 100 km/hr

37. i. In $\triangle APD$ and $\triangle BQC$

$AD = BC$ (given)

$AP = CQ$ (opposite sides of rectangle)

$\angle APD = \angle BQC = 90^\circ$

By RHS criteria $\triangle APD \cong \triangle CQB$

ii. $\triangle APD \cong \triangle CQB$

Corresponding part of congruent triangle

side $PD =$ side BQ

iii. In $\triangle ABC$ and $\triangle CDA$

$AB = CD$ (given)

$BC = AD$ (given)

$AC = AC$ (common)

By SSS criteria $\triangle ABC \cong \triangle CDA$

OR

In $\triangle APD$

$$\angle APD + \angle PAD + \angle ADP = 180^\circ$$

$$\Rightarrow 90^\circ + (180^\circ - 110^\circ) + \angle ADP = 180^\circ \text{ (angle sum property of } \triangle)$$

$$\Rightarrow \angle ADP = m = 180^\circ - 90^\circ - 70^\circ = 20^\circ$$

$$\angle ADP = m = 20^\circ$$

38. i. (c) 180°

ii. Show that in a right triangle the sum of legs is longest for an isosceles right triangle when hypotenuse remains same.

Take for example the length of diameter (hypotenuse) = 5 units.

Road D and Road B are equal hence (Road D = 3.53 units).

Let Road E be = 1, Road F = 4.89 units.

Therefore, length of Road B + Road D is greater than Road E + Road F.

iii. (c) Road G divides Road F into two equal.

iv. Yes, Priya is correct because arc corresponding to two equal roads (chords) are congruent.